

Biquandles with structures related to virtual links and twisted links

Naoko Kamada

Graduate School of Natural Sciences, Nagoya City University,
Aichi, 467-8501, Japan
kamada@nsc.nagoya-cu.ac.jp

and

Seiichi Kamada

Department of Mathematics, Hiroshima University,
Hiroshima 739-8526, Japan
kamada@math.sci.hiroshima-u.ac.jp

September 27, 2011

Abstract

We introduce two kinds of structures, called v-structures and t-structures, on biquandles. These structures are used for colorings of diagrams of virtual links and twisted links such that the numbers of colorings are invariants. Given a biquandle or a quandle, we give a method of constructing a biquandle with these structures. Using the numbers of colorings, we show that Bourgoin's twofoil and non-orientable virtual m -foils do not represent virtual links.

Keywords: Biquandles; virtual links; twisted links.

Mathematics Subject Classification 2000: 57M25

1 Introduction

A *virtual link diagram* is an oriented link diagram possibly with encircled crossings, called virtual crossings, that are neither positive crossings nor negative crossings. Two diagrams are *equivalent* if there is a sequence of the generalized Reidemeister moves defined in [16], which are generated by moves R1, $\dots$, R3, V1, $\dots$, V4 in Figure 1. The equivalence class of a virtual link diagram is called a *virtual link*. Virtual links correspond to stable equivalence classes of oriented links in the trivial I -bundles over closed orientable surfaces [5, 15, 19].

A *twisted link diagram* is a virtual link diagram which may have some bars on edges. Two diagrams are *equivalent* if there is a sequence of the extended Reidemeister moves defined in [3], which are generated by all moves in Figure 1. The equivalence class of a twisted link diagram is called a *twisted link*. (In [3] the extended Reidemeister moves are illustrated without orientations. Note that all moves with possible orientations are obtained by combining the moves in Figure 1. For example, see Figure 2.) Twisted links correspond to stable equivalence classes of oriented links in oriented 3-manifolds that are orientation I -bundles over closed but not necessarily orientable surfaces [3].

In this paper we define two kinds of structures on biquandles which are related to virtual links and twisted links.




Figure 1: The extended Reidemeister moves

A *biquandle* is a pair (X, R) consisting of a set X and a bijection $R : X^2 \rightarrow X^2$ satisfying certain conditions corresponding to Reidemeister moves for classical link diagrams [10, 12, 21] (Section 2). In Section 3 we introduce the notions of a v-structure $V : X^2 \rightarrow X^2$ and a t-structure $T : X \rightarrow X$ which are additional structures on a biquandle (X, R) related to virtual links and twisted links. The pair (V, T) is called a vt-structure of (X, R) .

A coloring of a virtual link diagram by a v-structured biquandle (X, R, V) or a coloring of a twisted link diagram by a vt-structured biquandle (X, R, V, T)




Figure 2: Another T1 move

is defined as follows: Let D be a diagram of a virtual link or a twisted link. The *edges* of D mean the connected arcs obtained when all the real crossings, virtual crossings and bars are removed.

Definition 1.1 A *coloring* of D by (X, R, V) or (X, R, V, T) is a map from the set of edges of D to X such that for each crossing or bar, say v , of D , if x_1, x_2, x_3, x_4 are elements of X assigned the edges around v as in Figure 3 then

- (1) $R(x_1, x_2) = (x_3, x_4)$ when v is a positive crossing,
- (2) $R^{-1}(x_1, x_2) = (x_3, x_4)$ when v is a negative crossing,
- (3) $V(x_1, x_2) = (x_3, x_4)$ when v is a virtual crossing, and
- (4) $T(x_1) = x_2$ when v is a bar.

We also call a coloring by (X, R, V) an (X, R, V) -*coloring*, and a coloring by (X, R, V, T) an (X, R, V, T) -*coloring*.

The concept of a coloring by (X, R, V) in Definition 1.1 and the following theorem (Theorem 1.2) were considered in [2]. Refer to [2] for examples.




Figure 3: Colorings

Theorem 1.2 ([2]) If D and D' are virtual link diagrams representing the same virtual link, then there is a bijection between the set of colorings of D by a v -structured biquandle (X, R, V) and that of D' .

This is generalized to twisted link diagrams.

Theorem 1.3 *If D and D' are twisted link diagrams representing the same twisted link, then there is a bijection between the set of colorings of D by a vt-structured biquandle (X, R, V, T) and that of D' .*

Therefore the number of colorings by a vt-structured biquandle (X, R, V, T) is an invariant of a twisted link.

Given a biquandle (X_0, R_0) and automorphisms f and g with $f^2 = 1$ and $fg = gf$, we give a method of constructing a vt-structured biquandle (X, R, V_f, T_g) , which we call a *twisted product* of (X_0, R_0) .

Let (X_0, R_0) be a biquandle. We use the notation due to [10] such that for $a, b \in X_0$,

$$R_0(a, b) = (b_a, a^b),$$

namely, $b_a = p_1 R_0(a, b)$ and $a^b = p_2 R_0(a, b)$, where $p_i : X_0 \times X_0 \rightarrow X_0$ is the i th factor projection.

Theorem 1.4 *Let (X_0, R_0) be a biquandle. Let $X = X_0 \times X_0$. Define a map $R : X^2 \rightarrow X^2$ by*

$$R((a_1, b_1), (a_2, b_2)) = ((a_{2a_1}, b_2^{b_1}), (a_1^{a_2}, b_{1b_2})).$$

For automorphisms f and g of (X_0, R_0) , define maps $V_f : X^2 \rightarrow X^2$ and $T_g : X \rightarrow X$ by

$$\begin{aligned} V_f((a_1, b_1), (a_2, b_2)) &= ((f^{-1}a_2, f^{-1}b_2), (fa_1, fb_1)), \text{ and} \\ T_g(a, b) &= (g^{-1}b, ga). \end{aligned}$$

Then the following holds.

- (1) (X, R) is a biquandle.
- (2) V_f is a v-structure of (X, R) .
- (3) Suppose that $f^2 = 1$ and $fg = gf$. Then (V_f, T_g) is a vt-structure of (X, R) .

Definition 1.5 In the situation of Theorem 1.4, we call the biquandle (X, R) the *twisted product biquandle* of (X_0, R_0) , and the vt-structured biquandle (X, R, V_f, T_g) a *twisted product* of (X_0, R_0) . When $f = g = 1$, we call the quadruplet the *standard twisted product* of (X_0, R_0) .

A *quandle* is a pair $(Q, *)$ consisting of a set Q and a binary operation $* : Q \times Q \rightarrow Q$, $(a, b) \mapsto a * b$, such that (i) for any $a \in Q$, $a * a = a$, (ii) for any $a, b \in Q$, there exists a unique element c with $c * b = a$, and (iii) for any $a, b, c \in Q$, $(a * b) * c = (a * c) * (b * c)$ [11, 14, 20]. The *dual* operation $\bar{*}$ of $*$ is a binary operation $\bar{*} : Q \times Q \rightarrow Q$ such that $a \bar{*} b = c \iff c * b = a$. In Section 4, we use Fenn and Rourke's notation [11]: $a * b$ and $a \bar{*} b$ are denoted by a^b and $a^{\bar{b}}$ (or $a^{b^{-1}}$), respectively, and a^{bc} means $(a^b)^c$, etc.

When (X_0, R_0) is the *biquandle derived from a quandle* $(Q, *)$, i.e., $X_0 = Q$ and $R_0(x, y) = (y, x * y)$, we call the twisted product biquandle (X, R) of (X_0, R_0) the *twisted product biquandle of $(Q, *)$* . If f and g are quandle automorphisms of $(Q, *)$, then they are biquandle automorphisms of (X_0, R_0) .

Suppose that $f^2 = 1$ and $fg = gf$. A *twisted product* of $(Q, *)$ means a twisted product (X, R, V_f, T_g) of (X_0, R_0) . The operations R, V_f, T_g on $X = Q^2$ are given as follows:

$$\begin{aligned} R((a_1, b_1), (a_2, b_2)) &= ((a_2, b_2 * b_1), (a_1 * a_2, b_1)), \\ V_f((a_1, b_1), (a_2, b_2)) &= ((fa_2, fb_2), (fa_1, fb_1)), \text{ and} \\ T_g(a, b) &= (g^{-1}b, ga). \end{aligned}$$

When $f = g = 1$, we have the following definition.

Definition 1.6 The *standard twisted product* of a quandle $(Q, *)$, which we denote by $\mathcal{B}(Q, *)$, is a vt-structured biquandle (X, R, V, T) such that $X = Q^2$ and

$$\begin{aligned} R((a_1, b_1), (a_2, b_2)) &= ((a_2, b_2 * b_1), (a_1 * a_2, b_1)), \\ V((a_1, b_1), (a_2, b_2)) &= ((a_2, b_2), (a_1, b_1)), \text{ and} \\ T(a, b) &= (b, a). \end{aligned}$$

For a quandle $(Q, *)$, the number of upper $(Q, *)$ -colorings of a virtual link diagram is an invariant of a virtual link, and so is that of lower $(Q, *)$ -colorings [16]. (A geometric interpretation of the upper/lower knot quandles of a virtual link is given in [15]. The upper/lower $(Q, *)$ -colorings correspond to the homomorphisms from these geometric quandles to $(Q, *)$ as in the classical case [11, 14, 20].)

Let $\mathcal{B}(Q, *)$ be the standard twisted product of a quandle $(Q, *)$.

Theorem 1.7 *If D is a twisted link diagram which is equivalent to a virtual link diagram D' , then the number of $\mathcal{B}(Q, *)$ -colorings of D is the product of the number of upper $(Q, *)$ -colorings of D' and that of lower $(Q, *)$ -colorings of D' .*

Corollary 1.8 *Let $(Q, *)$ be a finite quandle with n elements. If the number of $\mathcal{B}(Q, *)$ -colorings of a twisted link diagram D is less than n^2 , then D does not represent a virtual link.*

Proof. Every virtual link diagram has at least n trivial upper $(Q, *)$ -colorings and at least n trivial lower $(Q, *)$ -colorings. If D is equivalent to a virtual link diagram, by Theorem 1.7 there are at least n^2 colorings of D by $\mathcal{B}(Q, *)$. ■

Using an argument due to [8], Bourgoïn [3] showed that the twisted link diagram illustrated in Figure 4, which we call *Bourgoïn's twofoil*, does not represent a classical link. He also defined a twisted link invariant, called the *twisted Jones polynomial*, and showed that Bourgoïn's twofoil has the same twisted Jones polynomial with a certain virtual link diagram (Figures 6 and 7 of [3]). Thus, as mentioned in [3], one cannot distinguish it from virtual links by use of the twisted Jones polynomial.

We call a diagram on the left-hand side of Figure 5 a *non-orientable virtual m -foil* and denote it by F_m , where m is the number of the real crossings ($m \geq 1$). When $m = 2$, it is Bourgoïn's twofoil. It has a realization as a link diagram on




Figure 4: Bourgoin's twofoil

a projective plane depicted on the right-hand side of the figure, where the unit disk is made into a projective plane. By a calculation using induction on m , we see that the twisted Jones polynomial of F_m is

$$A^{-2m}(A^{-4m} + (-1)^{m+1}(1 + A^2 + A^{-2})) \in \mathbf{Z}[A^{\pm 1}, M].$$

This polynomial is also the twisted Jones polynomial of a virtual link diagram obtained when the two bars are removed from F_m . Thus one cannot distinguish it from virtual links by use of the twisted Jones polynomial.




Figure 5: non-orientable virtual m -foil

In Section 4 we study colorings of the diagram F_m by the standard twisted product $\mathcal{B}(Q, *)$, and show the following.

Theorem 1.9 (1) For $m > m' \geq 1$, F_m and $F_{m'}$ represent distinct twisted links.

(2) For $m \geq 1$, F_m does not represent a virtual link.

The first assertion of this theorem is also seen by the twisted Jones polynomials. Now we have an infinite family of twisted links which are not virtual links, but they are not distinguished from virtual links by the twisted Jones polynomials. This example was suggested the authors by Roger Fenn.

Remark 1.10 Given a quandle $(Q, *)$, let $X = Q^2$ and

$$\begin{aligned} R((a_1, b_1), (a_2, b_2)) &= ((a_2, b_2 \bar{*} b_1), (a_1 * a_2, b_1)), \\ V((a_1, b_1), (a_2, b_2)) &= ((a_2, b_2), (a_1, b_1)), \text{ and} \\ T(a, b) &= (b, a). \end{aligned}$$

Then (X, R) is a biquandle, and V is a v-structure of (X, R) . Since

$$\begin{aligned}(T \times T)R(T \times T)((a_1, b_1), (a_2, b_2)) &= ((a_2 \bar{*} a_1, b_2), (a_1, b_1 * b_2)) \quad \text{and} \\ VRV((a_1, b_1), (a_2, b_2)) &= ((a_2 * a_1, b_2), (a_1, b_1 \bar{*} b_2)),\end{aligned}$$

the operation $T : X^2 \rightarrow X^2$ does not satisfy $(T \times T)R(T \times T) = VRV$ unless $(Q, *)$ is an *involutory* quandle, i.e., $* = \bar{*}$. Thus the quadruplet (X, R, V, T) is not a vt-structured biquandle and one should not use this for colorings of twisted links.

We recall the notion of a biquandle, and prepare some lemmas in Section 2. In Section 3 the definitions of a t-structure and a v-structure are given, and Theorem 1.4 is proved. Section 4 is devoted to proofs of Theorems 1.3, 1.7 and 1.9.

2 Biquandles

For a set X we denote by X^n the n -fold Cartesian product of X , and denote by $p_i : X^n \rightarrow X$ the i th factor projection for each $i = 1, \dots, n$. The composition $g \circ f$ of two maps f and g is also denoted by $g \cdot f$ or gf . The identity map $x \mapsto x$ on X is denoted by 1_X or 1 , and the transposition map $(x, y) \mapsto (y, x)$ on X^2 is denoted by τ_{X^2} or τ .

The basic idea of a birack was given in [12]. The following is the definition of a (strong) birack and a (strong) biquandle introduced by R. Fenn, M. Jordan-Santana and L. Kauffman (Definitions 4.2 and 4.6 of [10]).

Definition 2.1 ([10]) A pair (X, R) of a set X and a bijection $R : X^2 \rightarrow X^2$ is a *birack* if the following conditions (B1) and (B2) are satisfied. It is a *biquandle* if (B1), (B2) and (B3) are satisfied.

(B1) R satisfies the set-theoretic Yang-Baxter equation, i.e.,

$$(R \times 1)(1 \times R)(R \times 1) = (1 \times R)(R \times 1)(1 \times R) : X^3 \rightarrow X^3.$$

(B2) For $a, b \in X$, let $f_a : X \rightarrow X$ and $f^b : X \rightarrow X$ be maps defined by

$$f_a(x) = p_1 R(a, x) \quad \text{and} \quad f^b(x) = p_2 R(x, b).$$

Then both f_a and f^b are bijections for every a and b of X .

(B3) For every a and b of X ,

$$(f_a)^{-1}(a) = f^{(f_a)^{-1}(a)}(a) \quad \text{and} \quad (f^b)^{-1}(b) = f_{(f^b)^{-1}(b)}(b).$$

As in Section 3 of [10] a birack/biquandle operation $R : X^2 \rightarrow X^2$ defines two binary operations on X ; $(a, b) \mapsto b_a$ and $(a, b) \mapsto a^b$ such that $R(a, b) = (b_a, a^b)$. Refer to [1, 4, 6, 9, 10, 13, 17] for examples of biracks and biquandles.

Let (X, R) and (X', R') be biracks or biquandles. A map $h : X \rightarrow X'$ is called a *homomorphism* if $(h \times h)R = R'(h \times h) : X^2 \rightarrow X'^2$. We denote it by $h : (X, R) \rightarrow (X', R')$. An *isomorphism* is a bijection which is a homomorphism.

Using the notion of a sideways operation due to Fenn, et al. [4, 9, 10], we can restate Definition 2.1 as follows.

Definition 2.2 A pair (X, R) consisting of a set X and a bijection $R : X^2 \rightarrow X^2$ is a *birack* if the following conditions (B1) and (B2') are satisfied. It is a *biquandle* if (B1), (B2') and (B3') are satisfied.

(B1) $(R \times 1)(1 \times R)(R \times 1) = (1 \times R)(R \times 1)(1 \times R) : X^3 \rightarrow X^3$.

(B2') There is a unique bijection $S : X^2 \rightarrow X^2$ such that for any $x_1, \dots, x_4 \in X$,

$$S(x_1, x_3) = (x_2, x_4) \iff R(x_1, x_2) = (x_3, x_4).$$

(B3') There is a bijection $s : X \rightarrow X$ such that for any $x \in X$,

$$R(x, s(x)) = (x, s(x)).$$

We call the bijections $S : X^2 \rightarrow X^2$ and $s : X \rightarrow X$ above the *sideways operation* of R and the *shift operation* of R , and denote them by $\text{side}R$ and $\text{shift}R$, respectively. (Note that if (B2') holds then a bijection $s : X \rightarrow X$ in (B3') is unique if there exists, since S is unique.) Refer to [4, 9, 10] for sideways operations.

First we observe that Definitions 2.1 and 2.2 are equivalent, and give some lemmas on biquandles.

Theorem 2.3 (1) For a bijection $R : X^2 \rightarrow X^2$, the conditions (B2) and (B2') are equivalent. (2) For a bijection $R : X^2 \rightarrow X^2$ satisfying (B2), the conditions (B3) and (B3') are equivalent.

Proof. (1) Suppose (B2). Define maps $S : X^2 \rightarrow X^2$ and $S^{-1} : X^2 \rightarrow X^2$ by

$$\begin{aligned} S(x, y) &= ((f_x)^{-1}(y), p_2 R(x, (f_x)^{-1}(y))) \quad \text{and} \\ S^{-1}(x, y) &= ((f^x)^{-1}(y), p_1 R((f^x)^{-1}(y), x)). \end{aligned}$$

Then $SS^{-1} = S^{-1}S = 1$, and $S(x_1, x_3) = (x_2, x_4) \iff R(x_1, x_2) = (x_3, x_4)$. Let $S' : X^2 \rightarrow X^2$ be another bijection such that $S'(x_1, x_3) = (x_2, x_4) \iff R(x_1, x_2) = (x_3, x_4)$. For any $x, y \in X$, since f_x is bijective, we have $p_1 S(x, y) = p_1 S'(x, y)$. Then $p_2 S(x, y) = p_2 R(x, p_1 S(x, y)) = p_2 R(x, p_1 S'(x, y)) = p_2 S'(x, y)$. Thus $S = S'$.

Suppose (B2'). The inverse maps of f_a and f^b are obtained by

$$(f_a)^{-1}(x) = p_1 S(x, a) \quad \text{and} \quad (f^b)^{-1}(x) = p_1 S^{-1}(b, x).$$

(2) Suppose (B3). Let $s : X \rightarrow X$ and $s^{-1} : X \rightarrow X$ be maps defined by

$$s(x) = (f_x)^{-1}(x) \quad \text{and} \quad s^{-1}(y) = (f^y)^{-1}(y).$$

Since $s(x) = (f_x)^{-1}(x) = f^{(f_x)^{-1}(x)}(x) = f^{s(x)}(x)$, we have $R(x, s(x)) = (x, s(x))$. Since $s^{-1}(y) = (f^y)^{-1}(y) = f^{(f^y)^{-1}(y)}(y) = f^{s^{-1}(y)}(y)$, we have $R(s^{-1}(y), y) = (s^{-1}(y), y)$. Then $s : X \rightarrow X$ and $s^{-1} : X \rightarrow X$ are the inverse maps of each other.

Suppose (B3'). Since $R(a, s(a)) = (a, s(a))$, we have $(f_a)^{-1}(a) = s(a) = f^{(f_a)^{-1}(a)}(a)$. Since $R(s^{-1}(b), b) = (s^{-1}(b), b)$, we have $(f^b)^{-1}(b) = s^{-1}(b) = f^{(f^b)^{-1}(b)}(b)$. ■

Now one may use Definition 2.2 instead of Definition 2.1.

We give some lemmas on biquandles, which are also valid for biracks.

Lemma 2.4 (1) *If (X, R) is a biquandle, then (X, R^{-1}) is a biquandle with $\text{side}R^{-1} = \tau \cdot \text{side}R \cdot \tau$ and $\text{shift}R^{-1} = \text{shift}R$.*

Proof. Since $R^{-1} \times 1 = (R \times 1)^{-1} : X^3 \rightarrow X^3$, the map R^{-1} satisfies the set-theoretic Yang-Baxter equation. Put $S' = \tau \cdot \text{side}R \cdot \tau$. Then $S'(x_3, x_1) = (x_4, x_2) \iff \text{side}R(x_1, x_3) = (x_2, x_4) \iff R(x_1, x_2) = (x_3, x_4) \iff (x_1, x_2) = R^{-1}(x_3, x_4)$. The uniqueness of S' follows from that of $\text{side}R$. Let $s = \text{shift}R$. Since $R(x, s(x)) = (x, s(x))$, we have $R^{-1}(x, s(x)) = (x, s(x))$. ■

Lemma 2.5 *If (X, R) is a biquandle, then $(X, \tau R \tau)$ is a biquandle with $\text{side}(\tau R \tau) = (\text{side}R)^{-1}$ and $\text{shift}(\tau R \tau) = (\text{shift}R)^{-1}$.*

Proof. It is obvious that $\tau R \tau$ is bijective. For simplicity we denote by τ_1, τ_2, R_1 and R_2 the maps $\tau \times 1, 1 \times \tau, R \times 1$ and $1 \times R$, respectively. Noting that $\tau_1 \tau_2 \tau_1 = \tau_2 \tau_1 \tau_2$ and $R_1 \tau_2 \tau_1 = \tau_2 \tau_1 R_2$, we see that $(\tau_1 R_1 \tau_1)(\tau_2 R_2 \tau_2)(\tau_1 R_1 \tau_1) = \tau_1 \tau_2 \tau_1 R_1 R_2 R_1 \tau_1 \tau_2 \tau_1$ and $(\tau_2 R_2 \tau_2)(\tau_1 R_1 \tau_1)(\tau_2 R_2 \tau_2) = \tau_2 \tau_1 \tau_2 R_2 R_1 R_2 \tau_2 \tau_1 \tau_2$. Thus $R_1 R_2 R_1 = R_2 R_1 R_2$ implies that

$$(\tau_1 R_1 \tau_1)(\tau_2 R_2 \tau_2)(\tau_1 R_1 \tau_1) = (\tau_2 R_2 \tau_2)(\tau_1 R_1 \tau_1)(\tau_2 R_2 \tau_2).$$

So $\tau R \tau$ satisfies the set-theoretic Yang-Baxter equation. $\tau R \tau(x_1, x_2) = (x_3, x_4) \iff R(x_2, x_1) = (x_4, x_3) \iff \text{side}R(x_2, x_4) = (x_1, x_3) \iff (\text{side}R)^{-1}(x_1, x_3) = (x_2, x_4)$. Thus $\text{side}(\tau R \tau) = (\text{side}R)^{-1}$. (The uniqueness of $\text{side}(\tau R \tau)$ follows from that of $\text{side}R$.) Let $s = \text{shift}R$. Since $R(x, s(x)) = (x, s(x))$ for every $x \in X$, we have $\tau T \tau(s(x), x) = (s(x), x)$. Thus $\text{shift}(\tau R \tau) = (\text{shift}R)^{-1}$. ■

Lemma 2.6 *Let (X, R) and (X', R') be biquandles, and let $f : X \rightarrow X'$ be a map. The following three conditions are mutually equivalent.*

- (i) *f is a homomorphism from (X, R) to (X', R') .*
- (ii) *f is a homomorphism from (X, R^{-1}) to (X', R'^{-1}) .*
- (iii) *f is a homomorphism from $(X, \tau R \tau)$ to $(X', \tau R' \tau)$.*

Proof. Since $(f \times f)R = R'(f \times f) \iff R'^{-1}(f \times f) = (f \times f)R^{-1}$, we have (i) $\iff$ (ii). Since $(f \times f)R = R'(f \times f) \iff \tau(f \times f)R\tau = \tau(f \times f)R'\tau \iff (f \times f)\tau R \tau = (f \times f)\tau R' \tau$, we have (i) $\iff$ (iii). ■

Lemma 2.7 *Let (X_1, R_1) and (X_2, R_2) be biquandles. Let $R : (X_1 \times X_2)^2 \rightarrow (X_1 \times X_2)^2$ be a map defined by*

$$R((a_1, b_1), (a_2, b_2)) = ((p_1 R_1(a_1, a_2), p_1 R_2(b_1, b_2)), (p_2 R_1(a_1, a_2), p_2 R_2(b_1, b_2))).$$

Then $(X_1 \times X_2, R)$ is a biquandle. Moreover, if $f_i : (X_i, R_i) \rightarrow (X_i, R_i)$ ($i = 1, 2$) are homomorphisms, then $f_1 \times f_2 : (X_1 \times X_2, R) \rightarrow (X_1 \times X_2, R)$ is a homomorphism.

Proof. The operation R satisfies the set-theoretic Yang-Baxter equation, since R_1 and R_2 do. The sideways operation S of R is given by

$$S((a_1, b_1), (a_3, b_3)) = ((p_1 S_1(a_1, a_3), p_1 S_2(b_1, b_3)), (p_2 S_1(a_1, a_3), p_2 S_2(b_1, b_3))),$$

where S_1 and S_2 are the sideways opetations of R_1 and R_2 . The shift operation s of R is given by

$$s((a, b)) = (s_1(a), s_2(b)),$$

where s_1 and s_2 are the shift opetations of R_1 and R_2 . It is obvious that $f_1 \times f_2$ is a homomorphism. ■

We call the biquandle $(X_1 \times X_2, R)$ in Lemma 2.7 the *direct product* of (X_1, R_1) and (X_2, R_2) .

3 v- and t-Strucures on biquandles

First we introduce the notion of a v-structure.

Definition 3.1 (cf. [2]) Let (X, R) be a biquandle. A bijection $V : X^2 \rightarrow X^2$ is a *v-structure* of (X, R) if the following conditions are satisfied.

- (1) (X, V) is a biquandle.
- (2) $V^2 = 1 : X^2 \rightarrow X^2$.
- (3) $(V \times 1)(1 \times V)(R \times 1) = (1 \times R)(V \times 1)(1 \times V) : X^3 \rightarrow X^3$.

We call (X, R, V) a *v-structured biquandle*.

A v-structure is used for colorings of virtual link diagrams.

Example 3.2 Let (X, R) be a biquandle. Let V be the transposition $\tau : X^2 \rightarrow X^2$, $(x, y) \mapsto (y, x)$. It is a v-structure. A biquandle with this v-structure is used for colorings of virtual link diagrams in [6, 10, 13, 18], etc.

Example 3.3 Let (X, R) be a biquandle. Let $f : (X, R) \rightarrow (X, R)$ be an automorphism. Let $V : X^2 \rightarrow X^2$ be a map defined by $V(x_1, x_2) = (f^{-1}x_2, fx_1)$. It is a v-structure. A biquandle with this structure is called a *virtual biquandle*. See [17] and Definition 3.3 of [7].

We introduce the notion of a t-structure, or a vt-structure, which is related to twisted links.

Definition 3.4 Let (X, R, V) be a v-structured biquandle. A bijection $T : X \rightarrow X$ is a *t-structure* of (X, R, V) if the following conditions are satisfied.

- (1) $T^2 = 1$.
- (2) $V(T \times 1) = (1 \times T)V$.
- (3) $(T \times T)R(T \times T) = VRV$.

We call (X, R, V, T) a *vt-structured biquandle*, and (V, T) a *vt-structure* of (X, R) .

Since $V^2 = 1$, the condition (2) of Definition 3.4 is equivalent to that $V(1 \times T) = (T \times 1)V$.

Proof of Theorem 1.4. (1) By Lemma 2.5, when (X_0, R_0) is a biquandle, so is $(X_0, \tau R_0 \tau)$. By Lemma 2.7, (X, R) is a biquandle, since it is the direct product of (X_0, R_0) and $(X_0, \tau R_0 \tau)$. (2) It follows from Lemmas 2.6 and 2.7

that $f \times f : X = X_0 \times X_0 \rightarrow X = X_0 \times X_0$ is an automorphism of (X, R) . By Example 3.3 we have (2). The assertion (3) is verified by direct calculation, which is left to the reader. ■

4 Proof of Theorems

In this section we prove Theorems 1.3, 1.7 and 1.9.

Proof of Theorem 1.3. Let Δ be a 2-disk where a move in Figure 1 transforms D to D' . For each move, there is a bijection of the colorings of D and D' such that the corresponding colorings are the identical outside Δ . When the move is of type R1, R2 or R3, it follows from that (X, R) is a biquandle [10]. When the move is of type V1, V2 or V3, it follows from that (X, V) is a biquandle and $V^2 = 1$. The case of V4 follows from the equality $(V \times 1)(1 \times V)(R \times 1) = (1 \times R)(V \times 1)(1 \times V)$. The case of type T1 follows from $V(T \times 1) = (1 \times T)V$. The case of type T2 follows from $T^2 = 1$. The case of type T3 follows from $(T \times T)R(T \times R) = VRV$. ■

Proof of Theorem 1.7. Let $\mathcal{B}(Q, *) = (X, R, V, T)$. By Theorem 1.3 it is sufficient to show that for a virtual link diagram D there is a bijection between the set of (X, R, V, T) -colorings of D , denoted by $\text{Col}(D, (X, R, V, T))$, and the Cartesian product of $\text{Col}^u(D, (Q, *))$ and $\text{Col}^l(D, (Q, *))$, where $\text{Col}^u(D, (Q, *))$ is the set of upper $(Q, *)$ -colorings of D and $\text{Col}^l(D, (Q, *))$ is the set of lower $(Q, *)$ -colorings of D . Since D is a virtual link diagram, an (X, R, V, T) -coloring of D is nothing more than an (X, R, V) -coloring of D . Let $c : E(D) \rightarrow X$ be an (X, R, V) -coloring of D , where $E(D)$ is the set of edges of D . Let $c^u : E(D) \rightarrow Q$ and $c^l : E(D) \rightarrow Q$ be the maps defined by $c^u = p_1 c$ and $c^l = p_2 c$ where p_i is the i th factor projection $X = Q^2 \rightarrow Q$. Then c^u is an upper $(Q, *)$ -coloring of D , and c^l is a lower $(Q, *)$ -coloring of D . Conversely for any upper $(Q, *)$ -coloring c^u and any lower $(Q, *)$ -coloring c^l , the map $c = (c^u, c^l) : E(D) \rightarrow X$ is an (X, R, V) -coloring of D . ■

In order to prove Theorem 1.9, we give a proposition on $\mathcal{B}(Q, *)$ -colorings of the non-orientable virtual m -foil F_m .

Let $(Q, *)$ be a quandle and $\mathcal{B}(Q, *)$ the standard twisted quandle. Consider a $\mathcal{B}(Q, *)$ -coloring of the diagram F_m and let x, y, z and w be elements of Q^2 given the edges as depicted in Figure 5. Put $x = (x_1, x_2)$ and $y = (y_1, y_2) \in Q^2$.

Proposition 4.1 *In this situation, x_1 and y_1 satisfy*

$$x_1 = x_1^{(y_1 x_1)^m} \quad \text{and} \quad y_1 = y_1^{(x_1 y_1)^m}. \quad (1)$$

And x_2 and y_2 are determined by

$$\begin{cases} x_2 = y_1^{(x_1 y_1)^n} \\ y_2 = x_1^{(y_1 x_1)^{n-1} y_1} \end{cases} \quad \text{if } m = 2n, \text{ and } \begin{cases} x_2 = x_1^{(y_1 x_1)^n y_1} \\ y_2 = y_1^{(x_1 y_1)^n} \end{cases} \quad \text{if } m = 2n + 1. \quad (2)$$

Conversely, for any elements x_1 and y_1 of Q satisfying (1), there exists a unique $\mathcal{B}(Q, *)$ -coloring of F_m .

Proof. The standard twisted product of a quandle is given by

$$R((x_1, x_2), (y_1, y_2)) = ((y_1, y_2^{x_2}), (x_1^{y_1}, x_2)).$$

Applying that m times, we have

$$(z, w) = ((x_1^{(y_1 x_1)^{n-1} y_1}, x_2^{(y_2 x_2)^n}), (y_1^{(x_1 y_1)^n}, y_2^{(x_2 y_2)^{n-1} x_2})) \quad \text{if } m = 2n, \quad (3)$$

and

$$(z, w) = ((y_1^{(x_1 y_1)^n}, y_2^{(x_2 y_2)^n x_2}), (x_1^{(y_1 x_1)^n y_1}, x_2^{(y_2 x_2)^n})) \quad \text{if } m = 2n + 1. \quad (4)$$

Since $(x, y) = V(T \times T)(z, w)$, we have

$$\begin{cases} x_1 = y_2^{(x_2 y_2)^{n-1} x_2}, & x_2 = y_1^{(x_1 y_1)^n} \\ y_1 = x_2^{(y_2 x_2)^n}, & y_2 = x_1^{(y_1 x_1)^{n-1} y_1} \end{cases} \quad \text{if } m = 2n, \quad (5)$$

and

$$\begin{cases} x_1 = x_2^{(y_2 x_2)^n}, & x_2 = x_1^{(y_1 x_1)^n y_1} \\ y_1 = y_2^{(x_2 y_2)^n x_2}, & y_2 = y_1^{(x_1 y_1)^n} \end{cases} \quad \text{if } m = 2n + 1. \quad (6)$$

From (5) and (6) we have (2), and eliminating x_2 and y_2 we have (1). Conversely, for any elements x_1 and y_1 of Q satisfying (1), let x_2 and y_2 be as in (2). Then $x = (x_1, x_2)$ and $y = (y_1, y_2)$ satisfy (5) and (6). Thus we have a coloring. ■

For a quandle $(Q, *)$, let $\Delta_m(Q, *)$ denote a subset

$$\{(a, b) \in Q^2 \mid a = a^{(ba)^m}, b = b^{(ab)^m}\} \quad (7)$$

of Q^2 .

Corollary 4.2 *For any quandle $(Q, *)$, assigning (x_1, y_1) to a $\mathcal{B}(Q, *)$ -coloring of F_m as in Proposition 4.1 is a bijection from $\text{Col}(F_m, \mathcal{B}(Q, *))$, the set of $\mathcal{B}(Q, *)$ -colorings of F_m , to $\Delta_m(Q, *)$.*

Proof of Theorem 1.9. (1) Let $(Q_{2m}, *)$ be the dihedral quandle of order $2m$, i.e., $Q_{2m} = \mathbb{Z}/2m\mathbb{Z}$ and $a * b = 2b - a$. It is easily seen that $\Delta_m(Q_{2m}, *) = (Q_{2m})^2$. Thus, by Corollary 4.2, $\#\text{Col}(F_m, \mathcal{B}(Q_{2m}, *)) = \#\Delta_m(Q_{2m}, *) = 4m^2$. On the other hand, if $m > m' \geq 1$, then $\Delta_{m'}(Q_{2m}, *) \neq (Q_{2m})^2$, for $(1, 0) \in (Q_{2m})^2$ does not belong to $\Delta_{m'}(Q_{2m}, *)$. Hence, by Corollary 4.2 again, $\#\text{Col}(F_{m'}, \mathcal{B}(Q_{2m}, *)) = \#\Delta_{m'}(Q_{2m}, *) < 4m^2$. Therefore F_m and $F_{m'}$ represent distinct twisted link.

(2) Let $(Q_{2m+1}, *)$ be the dihedral quandle of order $2m + 1$. Since $(1, 0) \in (Q_{2m+1})^2$ does not belong to $\Delta_m(Q_{2m+1}, *)$, we have $\Delta_m(Q_{2m+1}, *) \neq (Q_{2m+1})^2$. By Corollary 4.2, $\#\text{Col}(F_m, \mathcal{B}(Q_{2m+1}, *)) = \#\Delta_m(Q_{2m+1}, *) < \#(Q_{2m+1})^2$. By Corollary 1.8, F_m does not represent a virtual link. ■

Acknowledgements. The authors would like to express their gratitude to Roger Fenn for his valuable suggestions. The first author is partially supported by JSPS KAKENHI 22540093 and Grant-in-Aid for Research in Nagoya City University. The second author is partially supported by JSPS KAKENHI 21340015.

References

- [1] A. Bartholomew and R. Fenn, *Quaternionic invariants of virtual knots and links*, J. Knot Theory Ramifications **17** (2008), 231–251.
- [2] A. Bartholomew and R. Fenn, *Biquandles of small size and some invariants of virtual and welded knots*, J. Knot Theory Ramifications **20** (2011), 943–954.
- [3] M. O. Bourgoin, *Twisted link theory*, Algebr. Geom. Topol. **8** (2008), 1249–1279.
- [4] S. Budden and R. Fenn, *The equation, $[B, (A - 1)(A, B)] = 0$ and virtual knots and links*, Fund. Math. **184** (2004), 19–29.
- [5] J. S. Carter, S. Kamada and M. Saito, *Stable equivalence of knots on surfaces and virtual knot cobordisms*, J. Knot Theory Ramifications **11** (2002), 311–322.
- [6] J. S. Carter, D. S. Silver, S. G. Williams, M. Elhamdadi and M. Saito, *Virtual knot invariants from group biquandles and their cocycles*, J. Knot Theory Ramifications **18** (2009), 957–972.
- [7] J. Cenicerós and S. Nelson, *Virtual Yang-Baxter cocycle invariants*, Trans. Amer. Math. Soc. **361** (2009), 5263–5283.
- [8] Y. V. Drobotukhina, *An analogue of the Jones polynomial for links in $\mathbf{RP}^3$ and a generalization of the Kauffman–Murasugi theorem*, Algebra i Analiz **2** (1990), 171–191.
- [9] R. Fenn, *Biquandles and their application to virtual knots and links*, J. Knot Theory Ramifications **18** (2009), 785–789.
- [10] R. Fenn, M. Jordan-Santana and L. Kauffman, *Biquandles and virtual links*, Topology Appl. **145** (2004), 157–175.
- [11] R. Fenn and C. Rourke, *Racks and links in codimension two*, J. Knot Theory Ramifications **1** (1992), 343–406.
- [12] R. Fenn, C. Rourke and B. Sanderson, *An introduction to species and the rack space*, In: Topics in Knot Theory, M. E. Bozhuyu (Ed.), Kluwer Academic, Dordrecht, 1993, pp. 33–55.
- [13] D. Hrencecin and L. H. Kauffman, *Biquandles for virtual knots*, J. Knot Theory Ramifications **16** (2007), 1361–1382.
- [14] D. Joyce, *A classifying invariants of knots, the knot quandle*, J. Pure Appl. Alg. **23** (1982), 37–65.
- [15] N. Kamada and S. Kamada, *Abstract link diagrams and virtual knots*, J. Knot Theory Ramifications **9** (2000), 93–106.
- [16] L. H. Kauffman, *Virtual knot theory*, European J. Combin. **20** (1999), 663–690.
- [17] L. H. Kauffman and V. O. Manturov, *Virtual biquandles*, Fund. Math. **188** (2005), 103–146.

- [18] L. H. Kauffman and D. E. Radford, *Bioriented quantum algebras and a generalized Alexander polynomial for virtual links*, in: Diagrammatic Morphisms and Application, Contemp. Math., **318**, Amer. Math. Soc., 2003, pp. 113–140.
- [19] G. Kuperberg, *What is a virtual link?*, Algebr. Geom. Topol. **3** (2003), 587–591.
- [20] S. Matveev, *Distributive groupoids in knot theory*, Mat. Sb. (N.S.) **119** (**161**) (1982) 78–88, 160; English translation: Math. USSR-Sb. **47** (1984), 73–83.
- [21] D. Stanovský, *On axioms of biquandles*, J. Knot Theory Ramifications **15** (2006), 931–933.

